



RCP del Adulto

REFERENCIAS

DESFIBRILACIÓN

- The Public Access Defibrillation Trial Investigators. Publicaccess defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:637-46.
- Valenzuela TD, Roe DJ, Nichol G, Clark LL, Spaite DW, Hardman RG. Outcomes of rapid defibrillation by security officers after cardiac arrest in casinos. *N Engl J Med* 2000;343:1206-9.
- Caffrey SL, Willoughby PJ, Pepe PE, Becker LB. Public use of automated external defibrillators. *N Engl J Med* 2002;347:1242-7.
- O'Rourke MF, Donaldson E, Geddes JS. An airline cardiac arrest program. *Circulation* 1997;96:2849-53.
- Page RL, Joglar JA, Kowal RC, et al. Use of automated external defibrillators by a U.S. airline. *N Engl J Med* 2000;343:1210-6.
- van Alem AP, Vrenken RH, de Vos R, Tijssen JG, Koster RW. Use of automated external defibrillator by first responders in out of hospital cardiac arrest: prospective controlled trial. *BMJ* 2003;327:1312.
- Myerburg RJ, Fenster J, Velez M, et al. Impact of community-wide police car deployment of automated external defibrillators on survival from out-of-hospital cardiac arrest. *Circulation* 2002;106:1058-64.
- Capucci A, Aschieri D, Piepoli MF, Bardy GH, Iconomu E, Arvedi M. Tripling survival from sudden cardiac arrest via early defibrillation without traditional education in cardiopulmonary resuscitation. *Circulation* 2002;106:1065-70.
- White RD, Bunch TJ, Hankins DG. Evolution of a communitywide early defibrillation programme experience over 13 years using police/fire personnel and paramedics as responders. *Resuscitation* 2005;65:279-83.
- Mosesso Jr VN, Davis EA, Auble TE, Paris PM, Yealy DM. Use of automated external defibrillators by police officers for treatment of out-of-hospital cardiac arrest. *Ann Emerg Med* 1998;32:200-7.
- Smith KL, McNeil JJ. Cardiac arrests treated by ambulance paramedics and fire fighters: The Emergency MedicalResponse Program. *Med J Aust* 2002;177:305-9.
- Kellermann AL, Hackman BB, Somes G, Kreth TK, Nail L, Dobyns P. Impact of first-responder defibrillation in an urban emergency medical services system. *JAMA* 1993;270:1708-13.
- Becker L, Eisenberg M, Fahrenbruch C, Cobb L. Public locations of cardiac arrest: implications for public access defibrillation. *Circulation* 1998;97:2106-9.
- Herlitz J, Bang A, Axelsson A, Graves JR, Lindqvist J. Experience with the use of automated external defibrillators in out of hospital cardiac arrest. *Resuscitation* 1998;37:3-7.
- Kellermann AL, Hackman BB, Dobyns P, Frazier C, Nail L. Engineering excellence: options to enhance firefighter compliance with standing orders for first-responder defibrillation. *Ann Emerg Med* 1993;22:1269-75.
- Macdonald RD, Swanson JM, Mottley JL, Weinstein C. Performance and error analysis of automated external defibrillator use in the out-of-hospital setting. *Ann Emerg Med* 2001;38:262-7.
- Sunde K, Eftestol T, Askenberg C, Steen PA. Quality assessment of defibrillation and advanced life support using data from the medical control module of the defibrillator. *Resuscitation* 1999;41:237-47.
- Cleland MJ, Maloney JP, Rowe BH. Problems associated with the Z-fold region of defibrillation electrodes. *J Emerg Med* 1998;16:157-61.



RCP del Adulto

- Davis EA, Mosesso Jr VN. Performance of police first responders in utilizing automated external defibrillation on victims of sudden cardiac arrest. *Prehosp Emerg Care* 1998;2:101-7.
- Ornato JP, Shipley J, Powell RG, Racht EM. Inappropriate electrical countershocks by an automated external defibrillator. *Ann Emerg Med* 1992;21:1278-82.
- Calle PA, Monsieurs KG, Buylaert WA. Unreliable post event report from an automated external defibrillator. *Resuscitation* 2001;50:357-61.
- Zafari AM, Zarter SK, Heggen V, et al. A program encouraging early defibrillation results in improved in-hospital resuscitation efficacy. *J Am Coll Cardiol* 2004;44:846-52.
- Berg RA, Hilwig RW, Kern KB, Sanders AB, Xavier LC, Ewy GA. Automated external defibrillation versus manual defibrillation for prolonged ventricular fibrillation: lethal delays of chest compressions before and after countershocks. *Ann Emerg Med* 2003;42:458-67.
- Domanovits H, Meron G, Sterz F, et al. Successful automatic external defibrillator operation by people trained only in basic life support in a simulated cardiac arrest situation. *Resuscitation* 1998;39:47-50.
- Cusnir H, Tongia R, Sheka KP, et al. In hospital cardiac arrest: a role for automatic defibrillation. *Resuscitation* 2004;63:183-8.
- Alp NJ, Rahman S, Bell JA, Shahi M. Randomised comparison of antero-lateral versus antero-posterior paddle positions for DC cardioversion of persistent atrial fibrillation. *Int J Cardiol* 2000;75:211-6.
- Mathew TP, Moore A, McIntyre M, et al. Randomised comparison of electrode positions for cardioversion of atrial fibrillation. *Heart* 1999;81:576-9.
- Garcia LA, Kerber RE. Transthoracic defibrillation: does electrode adhesive pad position alter transthoracic impedance? *Resuscitation* 1998;37:139-43.
- Kerber RE, Martins JB, Kelly KJ, et al. Self-adhesive preapplied electrode pads for defibrillation and cardioversion. *J Am Coll Cardiol* 1984;3:815-20.
- Botto GL, Politi A, Bonini W, Broffoni T, Bonatti R. External cardioversion of atrial fibrillation: role of paddle position on technical efficacy and energy requirements. *Heart* 1999;82:726-30.
- Kirchhof P, Eckardt L, Loh P, et al. Anterior-posterior versus anterior-lateral electrode positions for external cardioversion of atrial fibrillation: a randomised trial. *Lancet* 2002;360:1275-9.
- Kerber RE, Grayzel J, Hoyt R, Marcus M, Kennedy J. Transthoracic resistance in human defibrillation. Influence of body weight, chest size, serial shocks, paddle size and paddle contact pressure. *Circulation* 1981;63:676-82.
- Deakin CD, Sado DM, Petley GW, Clewlow F. Is the orientation of the apical defibrillation paddle of importance during manual external defibrillation? *Resuscitation* 2003;56:15-8.
- Pagan-Carlo LA, Spencer KT, Robertson CE, Dengler A, Birkett C, Kerber RE. Transthoracic defibrillation: importance of avoiding electrode placement directly on the female breast. *J Am Coll Cardiol* 1996;27:449-52.
- Kanz KG, Kay MV, Biberthaler P, et al. Susceptibility of automated external defibrillators to train overhead lines and metro third rails. *Resuscitation* 2004;62:189-98.
- Dalzell GW, Cunningham SR, Anderson J, Adgey AA. Electrode pad size, transthoracic impedance and success of external ventricular defibrillation. *Am J Cardiol* 1989;64:741-4.
- Thomas ED, Ewy GA, Dahl CF, Ewy MD. Effectiveness of direct current defibrillation: role of paddle electrode size. *Am Heart J* 1977;93:463-7.
- Samson RA, Atkins DL, Kerber RE. Optimal size of selfadhesive preapplied electrode pads in pediatric defibrillation. *Am J Cardiol* 1995;75:544-5.
- Atkins DL, Sirna S, Kieso R, Charbonnier F, Kerber RE. Pediatric defibrillation: importance of paddle size in determining transthoracic impedance. *Pediatrics* 1988;82:914-8.
- Atkins DL, Kerber RE. Pediatric defibrillation: current flow is improved by using "adult" electrode paddles. *Pediatrics* 1994;94:90-3.



RCP del Adulto

- Hoyt R, Grayzel J, Kerber RE. Determinants of intracardiac current in defibrillation. Experimental studies in dogs. *Circulation* 1981;64:818-23.
- Killingsworth CR, Melnick SB, Chapman FW, et al. Defibrillation threshold and cardiac responses using an external biphasic defibrillator with pediatric and adult adhesive patches in pediatric-sized piglets. *Resuscitation* 2002;55:177-85.
- Dahl CF, Ewy GA, Warner ED, Thomas ED. Myocardial necrosis from direct current countershock: effect of paddle electrode size and time interval between discharges. *Circulation* 1974;50:956-61.
- Deakin CD, McLaren RM, Petley GW, Clewlow F, Dalrymple-Hay MJ. A comparison of transthoracic impedance using standard defibrillation paddles and self-adhesive defibrillation pads. *Resuscitation* 1998;39:43-6.
- Kerber RE, Martins JB, Ferguson DW, et al. Experimental evaluation and initial clinical application of new self-adhesive defibrillation electrodes. *Int J Cardiol* 1985;8:57-66.
- Deakin CD. Paddle size in defibrillation. *Br J Anaesth* 1998;81:657-8.
- Kirchhof P, Monnig G, Wasmer K, et al. A trial of selfadhesive patch electrodes and hand-held paddle electrodes for external cardioversion of atrial fibrillation (MOBIPAPA). *Eur Heart J* 2005;26:1292-7.
- Bojar RM, Payne DD, Rastegar H, Diehl JT, Cleveland RJ. Use of self-adhesive external defibrillator pads for complex cardiac surgical procedures. *Ann Thorac Surg* 1988;46:587-8.
- Brown J, Rogers J, Soar J. Cardiac arrest during surgery and ventilation in the prone position: a case report and systematic review. *Resuscitation* 2001;50:233-8.
- Wilson RF, Sirna S, White CW, Kerber RE. Defibrillation of high-risk patients during coronary angiography using self-adhesive, preapplied electrode pads. *Am J Cardiol* 1987;60:380-2.
- Bradbury N, Hyde D, Nolan J. Reliability of ECG monitoring with a gel pad/paddle combination after defibrillation. *Resuscitation* 2000;44:203-6.
- Callaway CW, Sherman LD, Mosezzo Jr VN, Dietrich TJ, Holt E, Clarkson MC. Scaling exponent predicts defibrillation success for out-of-hospital ventricular fibrillation cardiac arrest. *Circulation* 2001;103:1656-61.
- Eftestol T, Sunde K, Aase SO, Husoy JH, Steen PA. Predicting outcome of defibrillation by spectral characterization and nonparametric classification of ventricular fibrillation in patients with out-of-hospital cardiac arrest. *Circulation* 2000;102:1523-9.
- Eftestol T, Wik L, Sunde K, Steen PA. Effects of cardiopulmonary resuscitation on predictors of ventricular fibrillation defibrillation success during out-of-hospital cardiac arrest. *Circulation* 2004;110:10-5.
- Weaver WD, Cobb LA, Dennis D, Ray R, Hallstrom AP, Copass MK. Amplitude of ventricular fibrillation waveform and outcome after cardiac arrest. *Ann Intern Med* 1985;102:53-5.
- Brown CG, Dzwonczyk R. Signal analysis of the human electrocardiogram during ventricular fibrillation: frequency and amplitude parameters as predictors of successful countershock. *Ann Emerg Med* 1996;27:184-8.
- Callahan M, Braun O, Valentine W, Clark DM, Zegans C. Prehospital cardiac arrest treated by urban first-responders: profile of patient response and prediction of outcome by ventricular fibrillation waveform. *Ann Emerg Med* 1993;22:1664-77.
- Strohmenger HU, Lindner KH, Brown CG. Analysis of the ventricular fibrillation ECG signal amplitude and frequency parameters as predictors of countershock success in humans. *Chest* 1997;111:584-9.
- Strohmenger HU, Eftestol T, Sunde K, et al. The predictive value of ventricular fibrillation electrocardiogram signal frequency and amplitude variables in patients with outof-hospital cardiac arrest. *Anesth Analg* 2001;93:1428-33.
- Podbregar M, Kovacic M, Podbregar-Mars A, Brezocnik M. Predicting defibrillation success by 'genetic' programming in patients with out-of-hospital cardiac arrest. *Resuscitation* 2003;57:153-9.



RCP del Adulto

- Monsieurs KG, De Cauwer H, Wuyts FL, Bossaert LL. A rule for early outcome classification of out-of-hospital cardiac arrest patients presenting with ventricular fibrillation. *Resuscitation* 1998;36:37-44.
- Menegazzi JJ, Callaway CW, Sherman LD, et al. Ventricular fibrillation scaling exponent can guide timing of defibrillation and other therapies. *Circulation* 2004;109:926-31.
- Povoas HP, Weil MH, Tang W, Bisera J, Klouche K, Barbatsis A. Predicting the success of defibrillation by electrocardiographic analysis. *Resuscitation* 2002;53:77-82.
- Noc M, Weil MH, Tang W, Sun S, Pernat A, Bisera J. Electrocardiographic prediction of the success of cardiac resuscitation. *Crit Care Med* 1999;27:708-14.
- Strohmenger HU, Lindner KH, Keller A, Lindner IM, Pfenninger EG. Spectral analysis of ventricular fibrillation and closed-chest cardiopulmonary resuscitation. *Resuscitation* 1996;33:155-61.
- Noc M, Weil MH, Gazmuri RJ, Sun S, Biscera J, Tang W. Ventricular fibrillation voltage as a monitor of the effectiveness of cardiopulmonary resuscitation. *J Lab Clin Med* 1994;124:421-6.
- Lightfoot CB, Nremt P, Callaway CW, et al. Dynamic nature of electrocardiographic waveform predicts rescue shock outcome in porcine ventricular fibrillation. *Ann Emerg Med* 2003;42:230-41.
- Marn-Pernat A, Weil MH, Tang W, Pernat A, Bisera J. Optimizing timing of ventricular defibrillation. *Crit Care Med* 2001;29:2360-5.
- Hamprecht FA, Achleitner U, Krismer AC, et al. Fibrillation power, an alternative method of ECG spectral analysis for prediction of countershock success in a porcine model of ventricular fibrillation. *Resuscitation* 2001;50:287-96.
- Amann A, Achleitner U, Antretter H, et al. Analysing ventricular fibrillation ECG-signals and predicting defibrillation success during cardiopulmonary resuscitation employing $M(\alpha)$ -histograms. *Resuscitation* 2001;50:77-85.
- Brown CG, Griffith RF, Van Ligten P, et al. Median frequency - a new parameter for predicting defibrillation success rate. *Ann Emerg Med* 1991;20:787-9.
- Amann A, Rheinberger K, Achleitner U, et al. The prediction of defibrillation outcome using a new combination of mean frequency and amplitude in porcine models of cardiac arrest. *Anesth Analg* 2002;95:716-22, table of contents.
- Morrison LJ, Dorian P, Long J, et al. Out-of-hospital cardiac arrest rectilinear biphasic to monophasic damped sine defibrillation waveforms with advanced life support intervention trial (ORBIT). *Resuscitation* 2005;66:149-57.
- van Alem AP, Chapman FW, Lank P, Hart AA, Koster RW. A prospective, randomised and blinded comparison of first shock success of monophasic and biphasic waveforms in out-of-hospital cardiac arrest. *Resuscitation* 2003;58:17-24.
- Schneider T, Martens PR, Paschen H, et al. Multicenter, randomized, controlled trial of 150-J biphasic shocks compared with 200- to 360-J monophasic shocks in the resuscitation of out-of-hospital cardiac arrest victims. Optimized Response to Cardiac Arrest (ORCA) Investigators. *Circulation* 2000;102:1780-7.
- Martens PR, Russell JK, Wolcke B, et al. Optimal Response to Cardiac Arrest study: defibrillation waveform effects. *Resuscitation* 2001;49:233-43.
- Stothert JC, Hatcher TS, Gupton CL, Love JE, Brewer JE. Rectilinear biphasic waveform defibrillation of out-of-hospital cardiac arrest. *Prehosp Emerg Care* 2004;8:388-92.
- Carpenter J, Rea TD, Murray JA, Kudenchuk PJ, Eisenberg MS. Defibrillation waveform and post-shock rhythm in outof-hospital ventricular fibrillation cardiac arrest. *Resuscitation* 2003;59:189-96.
- Faddy SC, Powell J, Craig JC. Biphasic and monophasic shocks for transthoracic defibrillation: A meta analysis of randomised controlled trials. *Resuscitation* 2003;58:9-16.
- Walsh SJ, McClelland AJ, Owens CG, et al. Efficacy of distinct energy delivery protocols comparing two biphasic defibrillators for cardiac arrest. *Am J Cardiol* 2004;94:378-80.



RCP del Adulto

- Gliner BE, Lyster TE, Dillion SM, Bardy GH. Transthoracic defibrillation of swine with monophasic and biphasic waveforms. *Circulation* 1995;92:1634-43.
- White RD, Russell JK. Refibrillation, resuscitation and survival in out-of-hospital sudden cardiac arrest victims treated with biphasic automated external defibrillators. *Resuscitation* 2002;55:17-23.
- Bain AC, Swerdlow CD, Love CJ, et al. Multicenter study of principles-based waveforms for external defibrillation. *Ann Emerg Med* 2001;37:5-12.
- Bardy GH, Gliner BE, Kudenchuk PJ, et al. Truncated biphasic pulses for transthoracic defibrillation. *Circulation* 1995;91:1768-74.
- Bardy GH, Marchlinski F, Sharma A, et al. Multicenter comparison of truncated biphasic shocks and standard damped sine wave monophasic shocks for transthoracic ventricular fibrillation. *Circulation* 1996;94:2507-14.
- Greene HL, DiMarco JP, Kudenchuk PJ, et al. Comparison of monophasic and biphasic defibrillating pulse waveforms for transthoracic cardioversion. *Biphasic Waveform Defibrillation Investigators. Am J Cardiol* 1995;75:1135-9.
- Higgins SL, Herre JM, Epstein AE, et al. A comparison of biphasic and monophasic shocks for external defibrillation. *Physio-Control Biphasic Investigators. Prehosp Emerg Care* 2000;4:305-13.
- Higgins SL, O'Grady SG, Banville I, et al. Efficacy of lower-energy biphasic shocks for transthoracic defibrillation: a follow-up clinical study. *Prehosp Emerg Care* 2004;8:262-7.
- Mittal S, Ayati S, Stein KM, et al. Comparison of a novel rectilinear biphasic waveform with a damped sine wave monophasic waveform for transthoracic ventricular defibrillation. *ZOLL Investigators. J Am Coll Cardiol* 1999;34:1595-601.
- Weaver WD, Cobb LA, Copass MK, Hallstrom AP. Ventricular defibrillation: a comparative trial using 175-J and 320-J shocks. *N Engl J Med* 1982;307:1101-6.
- Tang W, Weil MH, Sun S, et al. The effects of biphasic and conventional monophasic defibrillation on postresuscitation myocardial function. *J Am Coll Cardiol* 1999;34:815-22.
- Morgan JP, Hearne SF, Raizes GS, White RD, Giuliani ER. High-energy versus low-energy defibrillation: experience in patients (excluding those in the intensive care unit) at Mayo Clinic-affiliated hospitals. *Mayo Clin Proc* 1984;59:829-34.
- Joglar JA, Hamdan MH, Ramaswamy K, et al. Initial energy for elective external cardioversion of persistent atrial fibrillation. *Am J Cardiol* 2000;86:348-50.
- Gliner BE, Jorgenson DB, Poole JE, et al. Treatment of outof-hospital cardiac arrest with a low-energy impedancecompensating biphasic waveform automatic external defibrillator. *The LIFE Investigators. Biomed Instrum Technol* 1998;32:631-44.
- White RD, Blackwell TH, Russell JK, Snyder DE, Jorgenson DB. Transthoracic impedance does not affect defibrillation, resuscitation or survival in patients with out-of-hospital cardiac arrest treated with a non-escalating biphasic waveform defibrillator. *Resuscitation* 2005;64:63-9.